

What is Claimed is:  
~~claims~~

a

1. Method for detecting nucleic acids in a sample comprising the steps:
- (a) purifying the nucleic acids in a binding space during which the nucleic acids are immobilized and impurities are separated,
  - (b) eluting the immobilized nucleic acids,
  - (c) amplifying the purified nucleic acids in an amplification space and
  - (d) detecting the amplification products in a detection space

**wherein**

the amplification space contains at least a part of the binding space.

2. Method as claimed in claim 1,

**wherein**

the detection space contains at least a part of the amplification space or/and at least a part of the binding space.

3. Method as claimed in claim 1 or 2,

**wherein**

an at least partial capillary space is used as the binding space or/and amplification space.

4. Method as claimed in one of the previous claims,

**wherein**

nucleic acids are adsorbed to a glass surface in step (a).

00780206-020901

5. Method as claimed in one of the previous claims,  
**wherein**  
a solution is used for the elution in step (b)  
which contains all reagents required for the  
amplification.
6. Method as claimed in one of the previous claims,  
**wherein**  
the amplification space can be thermostatted.
7. Method as claimed in claim 6,  
**wherein**  
the amplification space is surrounded by a heatable  
metal layer.
8. Method as claimed in one of the previous claims,  
**wherein**  
samples containing nucleic acids are lysed in step  
(a) before purification of the nucleic acids.
9. Method as claimed in one of the previous claims,  
**wherein**  
the sample contains cells.
10. Method as claimed in claim 9,  
**wherein**  
the cells are bound to a polystyrene surface.
11. Method as claimed in one of the previous claims,  
**wherein**  
the purification of the nucleic acids, the  
amplification of the purified nucleic acids and the  
detection of the amplification products are carried

09780206-02004

the same reaction space.

is claimed in one of the previous

are carried out in a closed d

the method as claimed in one of

detect pathogens in biological

or detecting nucleic acids in a

or by a method as claimed in one

to 12, comprising:

nding space to purify nucleic a

h the nucleic acids are immobi

rities are separated,

mplification space to amplify

tection space to detect nucleic

onally

ervoirs or/and supply lines for

nd reagents,

fication space contains at least

ng space.

claimed in claim 14,

tion space contains at least a

tion space or/and the binding s

claimed in claim 14 or 15,

ng space or/and amplification s

tially in the form of a capilla

12. Method as claimed in one of the previous claims,  
**wherein**  
all steps are carried out in a closed device.
13. Use of the method as claimed in one of the claims 1  
to 12 to detect pathogens in biological samples.
14. Device for detecting nucleic acids in a sample, in  
particular by a method as claimed in one of the  
claims 1 to 12, comprising:
- (a) a binding space to purify nucleic acids, in  
which the nucleic acids are immobilized and  
impurities are separated,
  - (b) an amplification space to amplify nucleic acids,
  - (c) a detection space to detect nucleic acids and  
optionally
  - (d) reservoirs or/and supply lines for the sample  
or/and reagents,
- wherein**  
the amplification space contains at least a part of  
the binding space.
15. Device as claimed in claim 14,  
**wherein**  
the detection space contains at least a part of the  
amplification space or/and the binding space.
16. Device as claimed in claim 14 or 15,  
**wherein**  
the binding space or/and amplification space is at  
least partially in the form of a capillary space.

17. Method for lysing a matrix containing nucleic acids,  
**wherein**  
a lysis mixture containing the matrix containing nucleic acids and a lysis reagent is moved through a capillary space during which the matrix is disrupted and the nucleic acids contained therein are released.
18. Method as claimed in claim 17,  
**wherein**  
the matrix containing nucleic acids comprises cells or/and cell fractions.
19. Method as claimed in claim 17 or 18,  
**wherein**  
a lysis reagent is used which contains a lytic enzyme or/and a chaotropic substance.
20. Method as claimed in one of the claims 17 to 19,  
**wherein**  
the capillary space is a glass capillary or/and a polystyrene capillary.
21. Method as claimed in claim 20,  
**wherein**  
the capillary space is a capillary coated with boron silicate.
22. Method as claimed in one of the claims 17 to 21,  
**wherein**  
the sample is passed several times through the capillary space.

23. Method as claimed in one of the claims 17 to 22,  
**wherein**  
the volume ratio of lysis mixture to capillary space  
is larger than 10:1.
24. Method for isolating nucleic acids from  
microorganisms,  
**wherein**  
a sample containing microorganisms is contacted with  
a polystyrene surface under conditions in which the  
microorganisms bind to the polystyrene surface and  
other sample components are separated, and the  
nucleic acids are isolated from the microorganisms.
25. Method as claimed in claim 24,  
**wherein**  
a salt is additionally added to facilitate the  
binding of the microorganisms to the polystyrene  
surface.
26. Method as claimed in claim 24 or 25,  
**wherein**  
a polystyrene capillary is used.
27. Method as claimed in one of the claims 24 to 26,  
**wherein**  
the sample is passed several times over the  
polystyrene surface.
28. Method as claimed in one of the claims 24 to 27,  
**wherein**  
the microorganisms are Chlamydia.

- add A27